

Perinatal Risk Factors for Morbidity vs Mortality

Derek A. Chapman, Ph.D. (TN Dept. of Health)


Tina Stanton-Chapman, M.S.Ed. (Vanderbilt U.)

Richard C. Urbano, Ph.D. (TN Dept. of Health)

Paper Presented at the 34th Annual Gatlinburg
Conference On Research and Theory in Intellectual
and Developmental Disabilities, Charleston, SC

March 5, 2001

USES OF EPIDEMIOLOGY



Epidemiological Methods

- Focus on proportions of cases
 - Low incidence conditions
 - Policy decisions
- Separate risk to individual from risk to the population

Risk Factor	Individual-Risk	Population-Risk
<i>Rare (LBW)</i>	High	Low
<i>Common (Poverty)</i>	Low	High


Risk Factors

- *Risk factor* refers to any characteristic of a
 - Person
 - Place
 - Time
- Identifiable prior to the event
- Can be causal or a marker for other factors


Electronic Data Linkage

- ***Florida:*** 1989-1990 birth certificates linked to 1997-98 public school records
 - 245,787 FL students who were born in FL
 - 41,612 (17%) in special education, not gifted
- ***Tennessee:*** Birth/death certificate linked file for 1989-1990 TN births
 - 145,355 live births
 - 1,487 (1%) were infant deaths

Risk Ratio Calculation Example



Epidemiological Concept of Risk Analysis



Individual-level Risk: RR(95%CI)

Infant Mortality

<i>apgar <4</i>	131.4 (119.7-144.3)
<i>vlbw</i>	88.5 (79.7-98.4)
<i>apgar 4-6</i>	31.7 (27.8-36.1)
<i>pre-term</i>	17.4 (15.6-19.4)

EH/SED

<i>male</i>	4.2 (3.8-4.5)
<i>mom ed <12</i>	3.7 (3.4-4.1)
<i>dad ed <12</i>	3.2 (2.8-3.7)
<i>apgar <4</i>	3.1 (1.6-6.1)
<i>unmarried</i>	3.1 (2.8-3.2)


Mild MR

<i>apgar 4-6</i>	8.3 (6.0-11.6)
<i>vlbw</i>	7.2 (6.2-8.4)
<i>mom ed <12</i>	5.5 (4.9-6.2)
<i>dad ed <12</i>	4.4 (3.9-5.1)
<i>apgar <4</i>	4.1 (3.3-5.0)


Mod/Sev/Prof MR

<i>apgar <4</i>	19.5 (11.5-33.0)
<i>vlbw</i>	13.1 (10.3-16.7)
<i>apgar 4-6</i>	12.6 (10.0-15.9)
<i>congen. abn.</i>	10.6 (8.8-12.9)

Very Low Birth Weight



Newborn Abnormal Cond.



Population Attributable Fraction (PAF) %

- Pulls together information about:
 - *Risk* associated with exposure to a risk factor (RR)
 - *Prevalence* of exposure to a risk factor (P_e)
- Weights risk ratio by % of population that has experienced the risk factor

$$PAF \% = \frac{P_e(RR - 1)}{1 + P_e(RR - 1)} \times 100$$

Population-level Risk: PAF%

Infant Mortality

Pe	RR	Risk Factor	PAF%
11%	17.4	<i>Pre-term</i>	65%
2%	62.0	<i>Apgar < 7</i>	54%
2%	88.5	<i>VLBW</i>	51%
35%	3.2	<i>Labor/Del. Comp.</i>	44%
11%	6.5	<i>Newborn Cond.</i>	38%

EH/SED

Pe	RR	Risk Factor	PAF%
52%	4.2	<i>Male</i>	62%
33%	3.1	<i>Unmarried</i>	41%
27%	3.7	<i>Mom Ed <12</i>	35%
34%	3.2	<i>Dad Ed <12</i>	23%

Mild MR

Pe	RR	Risk Factor	PAF%
27%	5.5	<i>Mom Ed <12</i>	44%
33%	2.8	<i>Unmarried</i>	37%
20%	4.4	<i>Dad Ed <12</i>	29%
44%	2.5	<i>Dad Ed =12</i>	28%

Mod/Sev/Prof MR

Pe	RR	Risk Factor	PAF%
52%	1.4	<i>Male</i>	19%
27%	2.0	<i>Mom Ed <12</i>	19%
33%	1.7	<i>Unmarried</i>	18%
8%	3.4	<i>Pre-term</i>	16%

Pe = Prevalence of Risk Factor in Population

Summary

- Patterns of risk vary based on:
 - Outcome
 - Level of analysis
- Epidemiological approach useful for developmental researchers
 - Target high-risk populations
 - Estimate potential effects of an intervention
 - Form etiologic hypotheses which guide basic research into biomedical cause (if present)